

WHAT IS CLAIMED IS:

1. A method for causing the elimination of an active principle, R, from cells of a mammal which contain the active principle, R, which comprises administering to said mammal, a compound identical in nature to said active principle in a homeopathic product of the formula  $RxCH$ , in which R is the active principle and in which xCH is a homeopathic dilution of said active ingredient R to eliminate R from the cells to restore normal function to the perturbed pericellular transport systems.
2. The method of claim 1, characterized in that the compound to be eliminated is a mineral compound and that the effect of its elimination is to restore normal function to an ion channel.
3. The method of claim 1, characterized in that the compound to be eliminated is a poison or part of a poison.
4. The method according to claim 1 which is for treating diseases caused by a metabolic error consisting either of an intracellular accumulation of a compound of simple or complex formula, or of an intracellular deficit of such a compound.
5. The method according to claim 4, characterized in that the consequence of the metabolic error is the production of abnormal metabolises, especially abnormal antibodies such as autoantibodies.
6. The method according to claim 4, characterized in that the metabolic error is genetically determined.
7. The method according to claim 5, characterized in that the metabolic error is genetically determined.

8. The method of claim 1, wherein xCH is hahnemannian or korsakowian homeopathic dilution.

9. The method of claim 1, wherein x is 4, 5, 7, 15 or 30.

10. The method of claim 9, wherein the active principle to be eliminated is a mineral compound and normal function is restored to an ion channel.

11. The method of claim 11, wherein x is 4, 5, 7, 15 or 30 and the active principle to be eliminated is a ~~poison~~ or part of a poison.

12. The method of claim 1, wherein the active principle to be eliminated is a virus or a viral particle.

13. A method of treating a disease caused by metabolic error consisting either or an intracellular accumulation of a compound of simple or complex formula, or of an intracellular deficit of such a compound which comprises administering to cells containing such a compound R or in need of said compound R, a homeopathic product having a formula RxCH in which R is said compound and in which xCH is its homeopathic dilution to treat said disease caused by metabolic error consisting either of an intracellular accumulation of said compound of simple or complex formula, or of an intracellular deficit of such a compound.

14. The method of claim 13, characterized in that the consequence of the metabolic error is the production of abnormal metabolites.

15. The method of claim 14, wherein the abnormal metabolites are abnormal antibodies.

16. The method of claim 15, wherein the abnormal antibodies are autoantibodies.

17. The method of claim 16, wherein the metabolic error is genetically determined.

18. The method of claim 16, wherein x is 4, 5, 7, 15 or 30.

19. The method of claim 2, wherein the mineral compound is a potassium, calcium or antimony containing mineral compound.

20. The method of claim 2, wherein R is a mineral compound which is a potassium, calcium or antimony containing mineral compound.

21. Application of homeopathic compounds of the general formula  $RxCH$  to the elimination of an intracellular chemical substance R from the cell, when this elimination causes the restoration of normal function to the perturbed pericellular transport systems, in order to obtain drugs which, by causing this elimination from the cell are intended for restoring normal function to the perturbed pericellular transport systems.

22. The application of claim 21, wherein  $xCH$  is hahnemannian or korsakowian homeopathic dilution, R is a mineral compound which is a potassium, calcium or antimony containing mineral compound and wherein x is 4, 5, 7, 15 or 30.